



Client Information

<i>Client Name</i>	Generic Client
<i>Date of birth (age)</i>	8 February 2001 (20)

Assessment Information

<i>Assessment</i>	World Health Organisation Disability Assessment Schedule 2.0 - Self (WHODAS-self)
<i>Date administered</i>	27 November 2021
<i>Assessor</i>	Dr David Hegarty
<i>Time taken</i>	1 minutes 49 seconds


Results

	Score (0 to 100)	Percentile	Average Score (0 to 4)	Descriptor
Overall Disability	40	90.38	1.72	Moderate
Cognition	67	98.14	2.67	Severe
Mobility	20	82.66	0.8	Mild
Self-Care	38	90.38	1.5	Moderate
Getting Along	15	78.42	0.6	Mild
Life Activities	72	98.14	2.88	Severe
Participation	31	88.35	1.25	Mild

Interpretive Text

None available

Average Scores





Client Name	Generic Client
--------------------	----------------

Scoring and Interpretation Information

There are two scoring methods used for the WHODAS 2.0:

1. Score (and its percentile)
2. Average score (and its descriptor)

The first score is determined using “item-response-theory” (IRT), where it takes into account multiple levels of difficulty for each WHODAS 2.0 item (1-36). This type of scoring for WHODAS 2.0 allows for more fine-grained analyses that make use of the full information of the response categories for comparative analysis across populations or subpopulations. It takes the coding for each item response as “none”, “mild”, “moderate”, “severe” and “extreme” separately, and then summarises the score by differentially weighting the items and the levels of severity. Converting the summary score into a metric ranging from 0 to 100 (where 0 = no disability; 100 = full disability). A percentile is provided that allows for a comparison to a large sample (n = 1,431) from a wide variety of populations (general population, populations with physical problems, populations with mental or emotional problems, populations with alcohol and drug use problems) from over 21 countries (WHO, 2010). A percentile of 50 indicates that an individual is experiencing an average level of disability when compared to other members of the sample.

The average scores are comparable to the WHODAS 5-point scale, which allows the clinician to think of the individual’s disability in terms of none (0-0.49), mild (0.5-1.49), moderate (1.5-2.49), severe (2.5-3.49), or extreme (3.5-4). The average domain and general disability scores were found to be reliable, easy to use, and clinically useful to the clinicians in the DSM-5 Field Trials (APA, 2021). The average domain score is calculated by dividing the raw domain score by the number of items in the domain. The average general disability score is calculated by dividing the raw overall score by number of items in the measure (i.e., 36).

The two scoring methods are used for each of the 6 domains:

1. Cognition – Assesses communication and thinking activities; specific areas assessed include concentrating, remembering, problem solving, learning and communicating.
2. Mobility – Assesses activities such as standing, moving around inside the home, getting out of the home and walking a long distance.
3. Self-care – Assesses hygiene, dressing, eating and staying alone.
4. Getting along – Assesses interactions with other people and difficulties that might be encountered with this life domain due to a health condition; in this context, “other people” includes those known intimately or well (e.g. spouse or partner, family members or close friends) and those not known well (e.g. strangers).



Client Name	Generic Client
--------------------	----------------

Scoring and Interpretation Information (cont.)

5. Life activities – Assesses difficulty with day-to-day activities (i.e. those that people do on most days, including those associated with domestic responsibilities, leisure, work and school).
 6. Participation – Assesses social dimensions, such as community activities; barriers and hindrances in the world around the respondent; and problems with other issues, such as maintaining personal dignity.

Client Responses

		None	Mild	Moderate	Severe	Extreme or cannot do
1	Concentrating on doing something for ten minutes?	0	1	2	3	4
2	Remembering to do important things?	0	1	2	3	4
3	Analysing and finding solutions to problems in day-to-day life?	0	1	2	3	4
4	Learning a new task, for example, learning how to get a new place?	0	1	2	3	4
5	Generally understanding what people say?	0	1	2	3	4
6	Starting and maintaining a conversation?	0	1	2	3	4
7	Standing for long periods such as 30 minutes?	0	1	2	3	4
8	Standing up from sitting down?	0	1	2	3	4
9	Moving around inside your home?	0	1	2	3	4
10	Getting out of your home?	0	1	2	3	4
11	Walking a long distance such as a kilometre (or equivalent)?	0	1	2	3	4
12	Washing your whole body?	0	1	2	3	4
13	Getting dressed?	0	1	2	3	4
14	Eating?	0	1	2	3	4
15	Staying by yourself for a few days?	0	1	2	3	4



Client Name | Generic Client

Client Responses (cont.)

		None	Mild	Moderate	Severe	Extreme or cannot do
16	Dealing with people you do not know?	0	1	2	3	4
17	Maintaining a friendship?	0	1	2	3	4
18	Getting along with people who are close to you?	0	1	2	3	4
19	Making new friends?	0	1	2	3	4
20	Sexual activities?	0	1	2	3	4
21	Taking care of your household responsibilities?	1	2	3	4	5
22	Doing most important household tasks well?	0	1	2	3	4
23	Getting all the household work done that you needed to do?	0	1	2	3	4
24	Getting your household work done as quickly as needed?	0	1	2	3	4
25	<small>If you work (paid, non-paid, self-employed) or go to school, complete questions 25-28 below. Otherwise, skip to question 29.</small> Because of your health condition, in the past 30 days, how much difficulty did you have in: Your day-to-day work/school?	0	1	2	3	4
26	Doing your most important work/school tasks well?	1	2	3	4	5
27	Getting all the work done that you need to do?	1	2	3	4	5
28	Getting your work done as quickly as needed?	0	1	2	3	4
29	<small>In the past 30 days:</small> How much of a problem did you have in joining in community activities (for example, festivities, religious or other activities) in the same way as anyone else can?	0	1	2	3	4
30	How much of a problem did you have because of barriers or hindrances in the world around you?	0	1	2	3	4
31	How much of a problem did you have living with dignity because of the attitudes and actions of others?	0	1	2	3	4
32	How much time did you spend on your health condition, or its consequences?	0	1	2	3	4
33	How much have you been emotionally affected by your health condition?	0	1	2	3	4
34	How much has your health condition been a drain on the financial resources of you or your family?	0	1	2	3	4



Client Name	Generic Client
--------------------	----------------

Client Responses (cont.)

		None	Mild	Moderate	Severe	Extreme or cannot do
35	How much of a problem did your family have because of your health problems?	0	1	2	3	4
36	How much of a problem did you have in doing things by yourself for relaxation or pleasure?	0	1	2	3	4
37	Overall, in the past 30 days, how many days were these difficulties present?					
	30					
38	In the past 30 days, for how many days were you totally unable to carry out your usual activities or work because of any health condition?					
	15					
39	In the past 30 days, not counting the days that you were totally unable, for how many days did you cut back or reduce your usual activities or work because of any health condition?					
	15					