

Client Information

Client Name | Generic Client

Date of birth (age) 14 December 1980 (43)

Assessment Information

Assessment Institute for Basic Research - Modified Overt Aggression Scale (IBR-

MOAS)

Date administered | 29 May 2024

Assessor Dr David Hegarty

Time taken | 1 minutes 20 seconds


Results

	Score (0 - 12)		
Verbal aggression towards others	7		
Verbal aggression towards self 2			
Physical aggression against people	2		
Physical aggression against objects	8		
Physical aggression against self	3		

Clusters

	Average Score (0-3)
Overt Aggression	1.4
Self-Destructive	0.6

Subscale Scores





Client Name

Generic Client

Scoring and Interpretation Information

Scores for the five areas of aggression measured are presented. Scores range from 0 - 12. Higher scores are indicative of more frequent and/or severe aggressive behaviours. The five areas of aggression are:

- 1. Verbal aggression towards others (Items 1-4)
- 2. Verbal aggression towards self (Items 5-8)
- 3. Physical aggression against other people (Items 9-12)
- 4. Physical aggression against objects (Items 13-16)
- 5. Physical aggression against self (Items 17-20)

A graph is presented to compare the five areas when first administered and when administered more than once a graph is presented to easily demonstrate change over time.

Additionally, Cohen et al. (2010) identified two clusters that are made up of the five areas of aggression:

1. Overt Aggression cluster (composed of Physical aggression against objects, Physical aggression against other people, and Verbal Aggression Toward Others areas of aggression) 2. Self Destructive cluster (composed of Physical aggression against self and Verbal aggression toward self)

These are presented as an average score between 0 - 3, calculated by the sum of the items in each cluster divided by the number it items. Average scores allow for comparison between the two clusters to determine if aggression is more overt or self-destructive in nature and to tailor treatment appropriately.

If administered more than once, a graph will be presented to visualise change in these clusters over time.

Cilent	Response	S

·						
		Never	Rarely	Sometimes	Often	
1	Makes loud noises, shouts angrily, screams at others	0	1	2	3	
2	Yells mild personal insults (e.g., "You're stupid!")	0	1	2	3	
3	Curses viciously, uses foul language in anger, makes moderate threats to others	0	1	2	3	
4	Makes clear threats of violence toward others ("I'm going to kill you") or requests help to control self	0	1	2	3	



Client Name | Generic Client

Client Responses (cont.) Never Rarely Sometimes Often Shows frustration and anger with self by making loud noises, screaming, moaning, or whining Makes mild personal insults toward self (e.g., "I'm stupid!" "I'm not good!") Curses angrily at self, talks more negatively about self ("I'm very bad" "I'm evil") Makes clear threats of violence toward self ("I want to die" "I don't want to live" "I want to kill myself") Makes threatening gestures, swings at people, grabs at clothes Strikes, kicks, pushes, pulls others' hair (without injury to them) Attacks others, causing mild to moderate physical injury (bruises, sprain, welts) Attacks others, causing severe physical injury (broken bones, deep lacerations, internal injury) Slams door, scatters clothing, makes a mess in anger Throws objects down, kicks furniture without breaking it, tries to tear clothes, marks the wall in anger Breaks objects in anger, smashes windows in anger, rips clothes in anger Sets fires, throws objects dangerously in anger Picks or scratches skin, hits self, pulls hair (with no or minor injury Bangs head, hits fist into objects, throws self onto floor or into objects (hurts self but without serious injury) Small cuts or bruises, minor burns as a result of self-injury Mutilates self, makes deep cuts, bites that bleed, internal injury, fracture, loss of consciousness, loss of teeth